

Observational Astronomy

Problems: Set 4

1. What is the expected value of the Fried parameter at a wavelength of 5500 Å if the observed seeing is 0.6 arcsec? What is the corresponding value of r_0 at 1.6 microns in the infrared assuming Kolmogorov turbulence?
2. Which has a greater energy flux, 10 photons $\text{cm}^{-2} \text{s}^{-1}$ at 10 Å or 10^5 photons $\text{cm}^{-2} \text{s}^{-1}$ at 5000 Å?
3. A star has a measured *I*-band magnitude of 22.0. How many photons per second are detected from this star by the William Herschel Telescope on La Palma (4.2 m diameter), assuming that the telescope and imaging optics have a throughput of 60%, the detector has a quantum efficiency of 80%, the sky has a brightness of 20 magnitudes per square arcsec, and the seeing is 1 arcsec. You can use the following information, for Vega, which has an *I*-band magnitude of $m_I = 0.0$:

Filter	λ_{eff} (Å)	$\Delta\lambda$ (Å)	F (erg $\text{s}^{-1} \text{cm}^{-2} \text{Å}^{-1}$)
I	7980	1500	1.13×10^{-9}

Estimate the exposure time required to detect the star at a signal-to-noise ratio of 20.

4. What fraction of the photons in the *V* band of a bright star would be absorbed by the atmosphere if one were to observe the star at an airmass of 2.5, and at the zenith (airmass = 1)? Assume that the atmospheric extinction $k(\lambda)$ in the *V* band is $0.15 \text{ mag airmass}^{-1}$.
5. In making differential observations, explain why you should know the colors of the variable and comparison stars.
6. Find the resolving power of a grating needed to separate the sodium spectral lines D_1 and D_2 , which are at 5895.944 Å and 5889.977 Å. How many lines must the grating have to achieve this resolution in second order?