



OBSERVATIONAL ASTRONOMY

AUTUMN 2024

Lecture 10

Vitaly Neustroev

Photometry: ST Magnitudes

388

The ST magnitude system is defined such that an object with constant flux $F_\lambda = 3.63 \times 10^{-9} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$ will have magnitude $ST = 0$ in every filter. In general,

$$ST_{mag} = -2.5 \log F_\lambda - 21.1$$

We will not discuss this system anymore.

Bolometric magnitudes

389

- **Bolometric magnitudes:** this gives a magnitude corresponding to the total flux integrated over **all wavelengths**
- The calculations are expressed as the difference between the bolometric magnitude and observed magnitude. The difference is then known as the bolometric correction: $BC = m_{\text{bol}} - V$
- The XXIXth IAU General Assembly in Honolulu recommended zero points for the absolute and apparent bolometric magnitude scales:
 - ▣ Resolution B2 defines the zero point of the absolute bolometric magnitude scale such that a radiation source with $M_{\text{bol}}=0$ has luminosity $L_0=3.0128 \times 10^{28}$ W.
 - ▣ The zero point of the apparent bolometric magnitude scale ($m_{\text{bol}}=0$) corresponds to irradiance $F_{\text{bol}} = 2.518 \times 10^{-8}$ W m⁻². The zero points were chosen so that the nominal **solar** luminosity (3.828×10^{26} W) corresponds to $M_{\text{bol}}(\text{Sun}) = 4.74$.
 - ▣ The nominal total solar irradiance (1361 W m⁻²) corresponds approximately to apparent bolometric magnitude $m_{\text{bol}}(\text{Sun}) = -26.832$.

Standard Stars for Photometry (1)

390

- The primary standards for the UBV system are a set of 10 bright, naked eye stars of magnitude 2 to 5, known as the North Pole sequence – comprise stars within 2° of the North pole star. The magnitudes of these stars define the UBV colour system.
- Instead of using the primary standards directly, we use a series of secondary standard stars, or just standard stars, whose magnitudes have been carefully measured relative to the primary stars.
- For broadband optical work (UBVRI filter system) the standard stars used most frequently today are from the work of the astronomer [Arlo Landolt](#). Landolt has devoted many years to measuring a set of standard star magnitudes.

Standard Stars for Photometry (2)

391

- What makes a good standard star?
 - ▣ A standard star must not be variable!
 - ▣ Standard stars must be of a brightness that will not overwhelm the detector and telescope in use, but must be bright enough to give a good S/N in a short exposure. For very large telescopes, many of the Landolt stars are too bright.
 - ▣ Ideally, a set of stars very close together in the sky will cover a wide range of colours.
 - ▣ Standard stars should be located across the sky so that they span a wide range of airmass.

Colour indices (1)

392

- **Colour indices:** this is the difference between magnitudes at two separate wavelengths:

$$C_{BV} = B - V; C_{VR} = V - R, \text{ and so on.}$$

- International colour index (outdated, but can be found in the literature) based upon photographic and photovisual magnitudes:

$$m_p - m_{pv} = C = B - V - 0.11$$

Colour indices (2)

393

- The $B - V$ colour index is closely related to the spectral type with an almost linear relationship for main sequence stars.
- For most stars, the B and V regions are located on the long wavelength side of the maximum spectral intensity.
- If we assume that the effective wavelengths of the B and V filters are 4400 and 5500 Å, then using the Planck equation:

$$L_{\lambda}(T) = \frac{2 h c_0^2}{\lambda^5} \left[\exp \left(\frac{h c_0}{\lambda k_B T} \right) - 1 \right]^{-1}$$

we obtain:

$$B - V \approx -2.5 \log \left[3.05 \frac{\exp(2.617 \times 10^4/T)}{\exp(3.27 \times 10^4/T)} \right]$$

Colour indices (3)

394

- For $T < 10000$ K this is approximately

$$B - V \approx -2.5 \log \left[3.05 \frac{\exp(2.617 \times 10^4/T)}{\exp(3.27 \times 10^4/T)} \right] = -1.21 + \frac{7090}{T}$$

The magnitude scale is an arbitrary one.

For $T = 9600$ K (Vega temperature), $B - V = 0.0$,

but we have obtained ~ 0.5 . Using this correction, we get:

$$T = \frac{7090}{(B - V) + 0.74} \text{ K}$$

Colour excess and Interstellar absorption

395

- More distant stars are affected by interstellar absorption, and since this is strongly inversely dependent upon wavelength.
- The colour excess measure the degree to which the spectrum is reddened:

$$E_{U-B} = (U - B) - (U - B)_0$$

$$E_{B-V} = (B - V) - (B - V)_0$$

where the subscript 0 denotes **unreddened** quantities – intrinsic colour indices.

- In the optical spectrum, interstellar absorption varies with both wavelength and the distance like this semi-empirical relationship:

$$A_\lambda = 6.5 \times 10^{-10} / \lambda - 2.0 \times 10^{-4} \text{ mag pc}^{-1}$$

where λ is in nanometers

Photometry

396

- Simple UBV photometry for hot stars results in determinations of temperature, Balmer discontinuity, spectral type, and reddening. From the latter we can estimate distance.
- Thus, we have a very high return of information for a small amount of observational effort. This is why the relatively crude methods of wideband photometry is so popular.

Photometry

397

Effective wavelengths (for an A0 star like Vega), absolute fluxes (corresponding to zero magnitude) and zeropoint magnitudes for the UBVRIJHKL Johnson-Cousins system

Bessell et al.
(1998, A&A, 333, 231)

Band	λ_c (Å)	$f_{\nu 0}$	$f_{\lambda 0}$	zp(f_λ)	zp(f_ν)
U	3660	1.790	417.5	-0.152	0.770
B	4380	4.063	632.0	-0.602	-0.120
V	5450	3.636	363.1	0.000	0.000
R	6410	3.064	217.7	0.555	0.186
I	7980	2.416	112.6	1.271	0.444
J	12200	1.589	31.47	2.655	0.899
H	16300	1.021	11.38	3.760	1.379
K	21900	0.64	3.961	4.906	1.886
L	34500	0.285	0.708	6.775	2.765

$$f_\lambda \text{ (} 10^{-11} \text{ erg s}^{-1} \text{cm}^{-2} \text{Å}^{-1}\text{)}$$
$$\text{mag}_\lambda = -2.5 \log (f_\lambda) - 21.100 - \text{zp}(f_\lambda)$$

$$f_\nu \text{ (} 10^{-20} \text{ erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1} = 1000 \text{ Jy)}$$
$$\text{mag}_\nu = -2.5 \log (f_\nu) - 48.585 - \text{zp}(f_\nu)$$

Photometry: Fun with Units (1)

398

- **Why do we continue to use magnitudes?**
 - Historical reasons: astronomers have built up a vast literature of catalogues and measurements in the magnitude system.
 - The magnitude system is logarithmic, which turns the huge range in brightness ratios into a much smaller range in magnitude differences: the difference between the Sun and the faintest star visible to the naked eye is only 32 magnitudes.
 - Simplicity: Astronomers have figured out how to use magnitudes in some practical ways which turn out to be easier to compute than the corresponding brightness ratios.
- However, in general converting between different magnitude and photometric systems is difficult: conversion factors depend on the spectrum of each object.

Photometry: Fun with Units (2)

399

- Astronomers who study objects outside the optical wavelengths do not have any historical measurements to incorporate into their work.
- In those regimes, measurements are almost always quoted in "more rational" systems: units which are linear with intensity (rather than logarithmic) and which become larger for brighter objects:
 - $\text{erg s}^{-1}\text{cm}^{-2} \text{\AA}^{-1}$
 - $\text{erg s}^{-1}\text{cm}^{-2} \text{Hz}^{-1}$

 - $1 \text{ Jansky [Jy]} = 10^{-26} \text{ W m}^{-2} \text{ Hz}^{-1} = 10^{-23} \text{ erg s}^{-1}\text{cm}^{-2} \text{ Hz}^{-1}$
 $F_{\nu} [\text{Jy}] = 3.34 \times 10^4 \lambda^2 F_{\lambda} [\text{erg s}^{-1} \text{cm}^{-2} \text{\AA}^{-1}]$
 $F_{\lambda} [\text{erg s}^{-1} \text{cm}^{-2} \text{\AA}^{-1}] = 3.00 \times 10^{-5} \lambda^{-2} F_{\nu} [\text{Jy}]$

Photometry: Fun with Units (3)

400

- Fluxes for a $V = 0$ star of spectral type A0 V at 5450 \AA :
 - $f_{\lambda}^0 = 3.63 \times 10^{-9} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$, or
 - $\varphi_{\lambda}^0 = f_{\lambda}^0 / h\nu = 996 \text{ photons s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$

- Useful:
 - $1 \text{ Jy} = 1.51 \times 10^3 / \lambda \text{ photons s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$

 - $\Delta\lambda / \lambda = 0.15 \text{ (U)}, 0.22 \text{ (B)}, 0.16 \text{ (V)}, 0.23 \text{ (R)}, 0.19 \text{ (I)}$

Night Sky Brightnesses

401

Lunar Age (days)	U	B	V	R	I
0	22.0	22.7	21.8	20.9	19.9
3	21.5	22.4	21.7	20.8	19.9
7	19.9	21.6	21.4	20.6	19.7
10	18.5	20.7	20.7	20.3	19.5
14	17.0	19.5	20.0	19.9	19.2

Signal from the sky background is present in every pixel of the aperture. Because each instrument generally has a different pixel scale, the sky brightness is usually tabulated for a site in units of mag/arcsecond².